

## Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation *in vitro*

Jun-Ho Byun<sup>1</sup>, Eun-Ju Kang<sup>2</sup>, Seong-Cheol Park<sup>1</sup>, Dong-Ho Kang<sup>3</sup>,  
Mun-Jeong Choi<sup>1</sup>, Gyu-Jin Rho<sup>2</sup>, Bong-Wook Park<sup>1</sup>

<sup>1</sup>Department of Oral and Maxillofacial Surgery, School of Medicine and Institute of Health Science, Gyeongsang National University,

<sup>2</sup>OBS/Theriogenology and Biotechnology, College of Veterinary Medicine, Gyeongsang National University,

<sup>3</sup>Department of Neurosurgery, School of Medicine, Gyeongsang National University, Jinju, Korea

**Abstract** (J Korean Assoc Oral Maxillofac Surg 2012;38:343-53)

**Objectives:** This aim of this study was to effectively isolate mesenchymal stem cells (hSMSCs) from human submandibular skin tissues (termed hSMSCs) and evaluate their characteristics. These hSMSCs were then chemically induced to the neuronal lineage and analyzed for their neurogenic characteristics *in vitro*.

**Materials and Methods:** Submandibular skin tissues were harvested from four adult patients and cultured in stem cell media. Isolated hSMSCs were evaluated for their multipotency and other stem cell characteristics. These cells were differentiated into neuronal cells with a chemical induction protocol. During the neuronal induction of hSMSCs, morphological changes and the expression of neuron-specific proteins (by fluorescence-activated cell sorting [FACS]) were evaluated.

**Results:** The hSMSCs showed plate-adherence, fibroblast-like growth, expression of the stem-cell transcription factors Oct 4 and Nanog, and positive staining for mesenchymal stem cell (MSC) marker proteins (CD29, CD44, CD90, CD105, and vimentin) and a neural precursor marker (nestin). Moreover, the hSMSCs in this study were successfully differentiated into multiple mesenchymal lineages, including osteocytes, adipocytes, and chondrocytes. Neuron-like cell morphology and various neural markers were highly visible six hours after the neuronal induction of hSMSCs, but their neuron-like characteristics disappeared over time (24-48 hrs). Interestingly, when the chemical induction medium was changed to Dulbecco's Modified Eagle Medium (DMEM) supplemented with fetal bovine serum (FBS), the differentiated cells returned to their hSMSC morphology, and their cell number increased. These results indicate that chemically induced neuron-like cells should not be considered true nerve cells.

**Conclusion:** Isolated hSMSCs have MSC characteristics and express a neural precursor marker, suggesting that human skin is a source of stem cells. However, the *in vitro* chemical neuronal induction of hSMSC does not produce long-lasting nerve cells and more studies are required before their use in nerve-tissue transplants.

**Key words:** Skin, Mesenchymal stem cell, *In vitro* neuronal differentiation

[paper submitted 2012. 8. 10 / revised 2012. 11. 15 / accepted 2012. 11. 22]

### I. Introduction

Recently, many researchers have tried to regenerate nerve tissue with tissue engineering techniques. Multipotent or pluripotent stem cells, cultured Schwann cells, and isolated

neural stem cells have been transplanted into nerve defect sites to improve peripheral and central nerve functions<sup>1-3</sup>. Among them, mesenchymal stem cells (MSCs) have been the focus in improving nerve regeneration because of their capability to provide multi-lineage differentiations and self-renewal potential<sup>4</sup>. Bone marrow-derived MSCs (BMSCs) can trans-differentiate *in vitro* into Schwann cell-like cells, which produce remarkable *in vivo* nerve regeneration when transplanted into a peripheral nerve defect<sup>4-7</sup>. Note, however, that bone marrow aspiration sometimes requires invasive procedures, possibly inducing a range of complications in patients such as pain, hemorrhage, and fear. Therefore, more accessible tissues such as skin or fat are investigated

#### **Bong-Wook Park**

Department of Oral and Maxillofacial Surgery, School of Medicine and Institute of Health Science, Gyeongsang National University, 79, Gangnam-ro, Jinju 660-702, Korea

TEL: +82-55-750-8263 FAX: +82-55-761-7024

E-mail: parkbw@gnu.ac.kr

©This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

\*This research was supported by the Basic Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science, and Technology (2012-0472).

as alternative sources of adult stem cells for the tissue engineering technique nowadays.


Recently, skin has been considered a potential adult stem cell source. It is highly accessible, and enough autologous tissue could be easily obtained with minimal donor site complications. Moreover, skin is an abundant pluripotent, multipotent cell source with immune privilege and potential for self-replication<sup>8-10</sup>. Several researchers have demonstrated that there are several different types of stem cells - such as skin-derived precursors (SKPs), skin-derived mesenchymal stem cells (SMSCs), and epidermal stem cells -- in the dermis and epidermis of skins<sup>10-14</sup>. In the previous study, we isolated porcine skin-derived cells from the ear skin of miniature pigs and showed the multipotency and MSC characteristics<sup>15</sup>. The cells were isolated from the epidermis and dermis in serum-containing medium, and they proliferated adherently on the culture plate and expressed MSC-marker proteins. In this study, human SMSCs (hSMSCs) from submandibular skin were isolated and cultured, and *in vitro* differentiation into mesenchymal cells such as osteocytes, adipocytes, and chondrocytes was evaluated under specific induction media. In addition, isolated hSMSCs were characterized by evaluating the expression of various cell surface markers (CD29, CD44, CD90, CD105, vimentin, and nestin) and transcription factors (Oct 4, Nanog, and Sox 2). Finally,

to evaluate the efficiency of their neural differentiation, hSMSCs were differentiated into neural cells under the chemically neural induction protocol of MSCs<sup>16,17</sup>, and various neurogenic and angiogenic proteins were evaluated by immunocytochemistry (ICC).

## II. Materials and Methods

### 1. Isolation and culture of hSMSCs

Human facial skin samples were obtained from four patients (2 males and 2 females; 24-45 years old, average of 33.3 years) who had undergone head and neck surgery via the submandibular approach.(Fig. 1. A) All experiments were authorized by the Gyeongsang National University Hospital Ethics Committee, and the patients gave their informed consent to tissue donation. Fresh human skin samples were transported to the laboratory, and hSMSCs were isolated as previous protocol<sup>15</sup>. Briefly, all hairs and subcutaneous fat tissues were removed, and the samples were then cut into 1-3 mm<sup>2</sup> explants containing the epidermis and dermis. Skin explants were attached to the culture plates, and 2 mL of Dulbecco's Modified Eagle Medium (DMEM)/F12 (1 : 1) (Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS; Invitrogen), 10 ng/mL epidermal



**Fig. 1.** Isolation and primary culture of human skin-derived cells with serum-containing adherent cell culture method (scale bar=100  $\mu$ m). A. Harvested submandibular skin tissue. B-D. Culturing human skin-derived cells (hSDCs) on the 3rd (B), 7th (C), and 14th (D) day can be observed during primary culture (P0). B. Irregular and heterogeneous hSDCs isolated from a skin fragment (black shadow) in the primary culture plates. C. After 7 days of P0, proliferating irregularly shaped hSDCs were detected in the plates. D. After about 2 weeks of P0, plate-adherent, fibroblast-like homogeneous cells were detected in the culture plates.

Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation *in vitro*. J Korean Assoc Oral Maxillofac Surg 2012

growth factor (EGF; Sigma-Aldrich, St. Louis, MO, USA), 10 ng/mL basic fibroblast growth factor (bFGF; Sigma-Aldrich), 100 U/mL penicillin (Sigma-Aldrich), and 100 µg/mL streptomycin (Sigma-Aldrich) were added. The culture plates were incubated at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub> in air for 3 or 5 days. After removing the remaining skin fragments, the attached cells were expanded *in vitro*, with the culture medium changed twice a week. Once confluent, the cells were dissociated using 0.25% (w/v) trypsin-ethylenediaminetetraacetic acid (EDTA; Invitrogen) solution and pelleted at 500 × g for 5 min. The cells were then re-grown and incubated until passage 3.

## 2. Cell surface and intracellular markers analysis

The cell surface and intracellular markers of hSMSCs at passage 3 were analyzed using a flow cytometer (BD FACSCalibur; Becton Dickinson and Company, Franklin Lakes, NJ, USA) in triplicate. Briefly, cells that reached 90% confluence were harvested using 0.25% EDTA and washed twice in Dulbecco's phosphate buffered saline (DPBS; Invitrogen) supplemented with 10% FBS. The cells for detecting CD44, CD90, and CD105 were labeled directly with fluorescein isothiocyanate (FITC)-conjugated CD markers (rat anti-mouse CD44 [1 : 100, BD Pharmingen; BD Biosciences, Franklin Lake, NJ, USA], mouse anti-human CD90 [1 : 100, BD Pharmingen], and goat anti-mouse CD105 [1 : 100, BD Pharmingen]). The cells were fixed in 3.7% formaldehyde for an hour to analyze the levels of CD29, vimentin, and nestin. After washing with DPBS, the samples were labeled with primary antibodies (mouse

anti-human CD29 [1 : 100, BD Pharmingen], mouse anti-human vimentin [1 : 100, Sigma-Aldrich], and mouse anti-human nestin [1 : 100, BD Pharmingen]) for 45 min at 37°C, followed by labeling with the FITC-conjugated secondary goat anti-mouse antibody (1 : 100, BD Pharmingen) for an hour.

## 3. *In vitro* neuronal differentiation

Neuronal differentiation of hSMSCs was performed using a modified chemical neural induction protocol for BMSCs differentiation<sup>16,17</sup>. Briefly, when hSMSCs at passage 3 reached 70% confluence, the cells were transferred to a neuronal preinduction medium containing DMEM (Invitrogen) with 20% FBS (Invitrogen) and 10 ng/mL bFGF (Sigma-Aldrich) for 24 hr. The cells were washed with PBS and cultured in neuronal induction medium consisting of DMEM supplemented with 2% dimethylsulfoxide (DMSO), 200 µM butylated hydroxyanisole (BHA), 25 mM KCl, 2 mM Valproic acid, 10 µM Forskolin, 1 µM Hydrocortisol, 5 µg/mL Insulin, and 2 mM L-glutamine without FBS for up to 48 hr. The cells were fixed for ICC at 0 hr, 6 hr, 24 hr, and 48 hr of induction. All supplemented chemicals in the neural induction medium were manufactured by Sigma-Aldrich Company.

To compare the morphological changes of the neuronally differentiated cells, the chemically inductive medium of the control cells was changed to DMEM supplemented with 20% FBS after 24 hr of chemical induction, and their morphological changes were observed after an additional 24 hr.

**Table 1.** Primary antibodies for the immunocytochemical study of chemically neural induced cells

Antibody	Type	Company	Catalog number	Dilution
Nestin	Mouse monoclonal	Sigma-Aldrich Co., St. Louis, MO, USA	61658	1 : 200
S-100	Rabbit polyclonal	ThermoScientific, Rockford, IL, USA	RB9018	1 : 300
NF	Goat polyclonal	Santa Cruz, Santa Cruz, CA, USA	sc-16143	1 : 100
NGF	Rabbit polyclonal	Santa Cruz	sc-548	1 : 100
p75NGFR	Mouse monoclonal	Santa Cruz	sc-13577	1 : 100
trkA	Goat polyclonal	Santa Cruz	sc-20537	1 : 100
VEGF	Rabbit polyclonal	Santa Cruz	sc-152	1 : 100
VEGFR1	Mouse monoclonal	Abcam, Cambridge, UK	ab9540	1 : 100
VEGFR2	Rabbit polyclonal	Abcam	ab71772	1 : 100
β-tubulin	Goat polyclonal	Santa Cruz	sc-9935	1 : 100
MBP	Goat polyclonal	Santa Cruz	sc-13914	1 : 100
NeuN	Mouse polyclonal	Chemicon, Temecula, CA, USA	MAB377	1 : 100

(NF: neurofilament, NGF: nerve growth factor, trkA: tyrosine kinase receptor A, VEGF: endothelial cell growth factor, MBP: myelin basic protein, NeuN: neural-specific nuclear protein)

Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation *in vitro*. J Korean Assoc Oral Maxillofac Surg 2012

4. Immunocytochemical analysis of hSMSCs and *in vitro* neural induced cells

When the hSMSCs reached passage 3, they were rinsed with PBS and fixed in 4% neutral buffered formaldehyde for 30 min at room temperature. ICC for the transcription factors (Oct 4, Nanog, and Sox 2) was conducted. A 1 : 200 dilution of primary goat polyclonal anti-human Oct 3/4 (sc-8628; Santa Cruz, Santa Cruz, CA, USA), a 1 : 200 dilution of primary goat polyclonal anti-human Nanog (sc-30331, Santa Cruz), and a 1 : 200 dilution of primary rabbit polyclonal anti-human Sox 2 (sc-20088, Santa Cruz) were used to detect the expression of transcription factors.

In the neural induction medium, the differentiated cells were fixed at 0 hr (immediately after preinduction), 6 hr, 24 hr, and 48 hr after neural induction. Table 1 lists the primary antibodies used for the evaluation of neural differentiation. Dilutions (1 : 100) of FITC-conjugated donkey anti-goat polyclonal IgG (Jackson ImmunoResearch Laboratories Inc., West Grove, PA, USA), FITC-conjugated donkey anti-rabbit polyclonal IgG (711-095-152, Jackson ImmunoResearch Laboratories Inc.), and FITC-conjugated goat anti-mouse polyclonal IgG (115-096-003, Jackson ImmunoResearch Laboratories Inc.) were used as secondary antibodies.

Densitometric analyses of each immunostaining were performed using analySIS TS software (Olympus Soft Imaging Solution, Münster, Germany). For the evaluation of one antibody's expression, at least three slides were immunostained and statistically analyzed at each time point.

5. Reverse transcription-polymerase chain reaction (RT-PCR) analysis

The hSMSCs at passage 3 were evaluated by RT-PCR for the expression of transcription factors Oct 4, Sox 2, and Nanog. Table 2 lists the RT-PCR primers for the markers used in this study. The total RNA was extracted from the cultured cells using an RNeasy Mini Kit (Qiagen, Valencia, CA, USA). cDNA synthesis was performed for 30 min at 55°C using an Omniscript Reverse Transcription Kit (Qiagen) with oligo-dT primers. The cDNAs produced were used as template for PCR amplification. PCR was performed using Maxime PCR Premix (iNtRON Biotechnology, Seongnam, Korea) under the following conditions: pre-denaturation at 94°C for 3 min, followed by 34 cycles of denaturation at 94°C for 45 s, annealing at 60°C or 58°C for 30 s, elongation at 72°C for 45 s, and final extension at 72°C for 10 min using a Thermocycler (PTC-200; GMI, Anoka, MN, USA).

6. Statistical analysis

All values of the counted and calculated cell numbers and intensities of immunostainings of the *in vitro* neural differentiated cells were statistically analyzed by the Manova test, and independent grouping variables were compared using Bonferroni and SPSS software version 18.0 (SPSS Inc., Chicago, IL, USA). Data were expressed as mean±SD. Differences were considered to be significant when  $P < 0.05$ .

**Table 2.** RT-PCR primers used for evaluating transcription factors and osteogenic and adipogenic differentiations

Gene	Sequence of primer (5' - 3')	Amplification size (bp)	Temperature (°C)	Locous
GAPDH	F - GAGTCAACGGATTTGGTCGT R - TTGATTTTGGAGGGATCTCG	238	60	AB062273
Oct 4	F - GATCCTCGGACCTGGCTAAG R - GACTCCTGCTTCACCCTCAG	213	60	AM851115
Nanog	F - CAAAGGCAAACAACCCACTT R - TCTGGAACCAGGTCTTCACC	218	60	AB093576
Sox2	F - TCACGTACACTGCCCTGAAG R - TGCAACGGATTGTGTTGTTT	175	60	Z31560
Osteonectin	F - GTGCAGAGGAAACCGAAGAG R - AAGTGGCAGGAAGAGTCGAA	202	60	NM_003118
Osteocalcin	F - GGCAGCGAGGTAGTGAAGAG R - CTGGAGAGGAGCAGAACTGG	230	60	X53698
PPAR $\gamma$ 2	F - ACTGCGCTACAAATGCACAC R - TTGATGCCGAGAAAGGAGAT	248	60	AJ563369
Ap2	F - TACTGGGCCAGGAATTTGAC R - ATGCGAACTTCAGTCCAGGT	237	58	NM_001442

(RT-PCR: reverse transcription-polymerase chain reaction, GAPDH: glyceraldehyde 3-phosphate dehydrogenase)

Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation *in vitro*. J Korean Assoc Oral Maxillofac Surg 2012

### III. Results

#### 1. Cell isolation and culture

Both floated sphere-forming cells and plate-adherent, fibroblast-like cells were co-detected in the culture medium from the first day of primary culture. After 3 days of primary culture, most cells had shown plate-adherent growth in the gelatin-coated plates. Initially, the attached skin-derived cells showed heterogeneously irregular shapes, and partial colony formations were observed. Note, however, that homogeneously shaped and plate-adherent fibroblast-like cells were mainly detected at the end - about 2 weeks later - of the primary culture.(Figs. 1. B-D) These homogeneously shaped, plate-adherent fibroblast-like cells were allowed to proliferate, reaching passage 2 or 3.


#### 2. Expression of transcription factors determined by RT-PCR and cell surface markers measured by FACS analysis

After human skin-derived cells were cultured to passage 3, the expression of transcription factors such as Oct 4, Nanog, and Sox 2 was evaluated by ICC and RT-PCR. Oct 4 and Nanog were highly visible in cultured adult hSMSCs, whereas

Sox 2 was hardly observed by ICC or RT-PCR.(Figs. 2. A, 2. B) The hSMSCs at passage 3 were positive for typical MSC markers (CD29, CD44, CD90, CD105, and vimentin) based on FACS analysis. In addition, the neural precursor marker, nestin, was detected in the hSDCs.(Fig. 3) The results above demonstrate that human skin-derived cells in this study were multipotent MSCs exhibiting the characteristics of a neural precursor. Moreover, the cultured hSMSCs in this study were successfully differentiated into mesenchymal lineage cells, osteocytes, adipocytes, and chondrocytes in specific induction media.(Fig. 4. A) These differentiated cells from hSMSCs also showed specific osteogenic and adipogenic marker proteins by RT-PCR.(Fig. 4. B)


#### 3. *In vitro* neural induction of hSMSCs

Specific morphological change from hSMSCs was not observed after neural preinduction.(Fig. 5. A) When hSMSCs were neuronally differentiated using various chemical compounds under FBS-deprived conditions, most strongly resembled neurons and included the retraction of cell body and process elaboration, and they were observed after 6 hr of induction.(Fig. 5. B) As the neuronal induction time reached 24 hrs and 48 hrs, however, the differentiated neuron-like cells decreased in number, and their cell morphology




**Fig. 2.** Expression of early transcription factors Oct 4, Nanog, and Sox 2 by immunocytochemistry (A: scale bar=100  $\mu$ m) and reverse transcription-polymerase chain reaction (B) in human skin-derived cells (hSDCs) at passage 3. Positive expression of Oct 4 and Nanog, even though Sox 2 was hardly expressed, indicates that the hSDCs in this study are multipotential primitive cells. (GAPDH: glyceraldehyde 3-phosphate dehydrogenase)

*Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation in vitro. J Korean Assoc Oral Maxillofac Surg 2012*




**Fig. 3.** Fluorescence-activated cell sorting analysis of cultured human skin-derived cells. Skin-derived cells at passage 3 were positive for specific mesenchymal stem cell markers (CD29, CD44, CD90, CD105, and vimentin) and neural precursor cell marker (nestin). Open histograms represent staining with negative control, with the black histograms depicting the fluorescence intensity of each of the cell surface antibodies.

Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation in vitro. J Korean Assoc Oral Maxillofac Surg 2012



**Fig. 4.** Mesenchymal-lineage differentiations of human skin-derived mesenchymal stem cells (hSMSCs) into osteocytes (a, b), adipocytes (c), and chondrocytes (d) for 4 weeks (A: scale bar=100  $\mu$ m). A. In vitro differentiated cells showed positive staining in the specific staining methods. (a, b) Calcium deposits were observed on the cell surface by von Kossa (a) and Alizalin red (b) staining. (c) Lipid droplets were noted in the cytoplasm of cells by Oil red O staining. (d) Proteoglycans were confirmed on the cell surface using Alcian blue. B. Reverse transcription-polymerase chain reaction results for *in vitro* differentiated osteocytes and adipocytes from hSMSCs. (a) ON and OC were detected in osteogenic differentiated cells. (b) PPAR $\gamma$ 2 and aP2 were expressed in adipogenic differentiated cells. (OC: osteocalcin, ON: osteonectin, GAPDH: glyceraldehyde 3-phosphate dehydrogenase, aP2: adipocyte protein 2)

Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation in vitro. J Korean Assoc Oral Maxillofac Surg 2012



**Fig. 5.** The upper graph illustrates the schematic *in vitro* neural induction protocol used in this study. Cultured hSMSCs at passage 3 were preinduced for 24 hrs. The experimental group was neurally induced by a chemical protocol for 48 hr. In the control group, 24 hr after neural induction, the inductive medium was changed to DMEM supplemented with 20% FBS, and morphologic changes were then observed after an additional 24 hrs of media change. A-E. The microphotographs show the morphologic changes of hSMSCs after chemical neural induction (scale bar=100 μm). A. Immediately after neuronal preinduction (0 hr). There are no remarkable morphological changes compared to the original hSMSCs. B. Six hours (6 hr) after neural induction, the neuron-like cells exhibit peak activity. C, D. After the passage of neural induction time (24 and 48 hrs post-neural induction), the neuron-like cells decreased in number, and their shape deteriorated. E. In the control cells, neural differentiated cells returned to the original hSMSC morphology, and cell number increased 24 hrs after media change as DMEM with 20% FBS. F. The number of cells decreased with the passage of neural induction time, but the number increased after the inductive medium was changed. (hSMSCs: human skin-derived mesenchymal stem cells, bFGF: basic fibroblast growth factor, FBS: fetal bovine serum, DMEM: Dulbecco's Modified Eagle Medium, DMSO: dimethylsulfoxide, VPA: valproic acid, BHA: butylated hydroxyanisole)

*Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation in vitro. J Korean Assoc Oral Maxillofac Surg 2012*


deteriorated (Figs. 5. C, 5. D) In the control cells, the induction medium was changed to DMEM with 20% FBS after 24 hrs of neural induction, and the neural differentiated cells returned to the original shape of hSMSCs.(Fig. 5. E) In addition, the number of cells increased 24 hrs after medium change.(Fig. 5. F)

In the immunocytochemical studies of differentiated neuron-like cells, neuron- and angiogenesis-related proteins were highly expressed, peaking 6 hrs after neural induction. Note, however, that the expression intensity of all proteins decreased as induction time reached 24 hrs and 48 hrs. Interestingly, the neural precursor marker, nestin, was substantially expressed in the pre-induced cells (0 hr). In addition, the enhanced co-expression of the nerve growth factor (NGF) and its two receptors (p75NGFR and trkA) as well as the vascular endothelial growth factor (VEGF) and its two receptors (VEGFR1 and VEGFR2) was detected at the early stages of neurogenic induction (6 hrs). The decreased expression

of these proteins was then observed after 24 hr and 48 hr of neurogenic induction.(Fig. 6)

#### IV. Discussion

Among the skin-derived stem cells, SKPs - believed to be endogenous embryonic neural crest-derived precursor cells that persist into adulthood - have been studied most widely. SKPs originated with the dermis of the skin and formed floating spheres in the serum-free culturing conditions supplemented with various growth factors - including FGF-2, EGF, and B27 - after enzymatic digestion and cell dissociation of the skin. Moreover, they have distinct characteristics from MSCs, and their full potential is normally restricted by the local environment, but such can be revealed under culture conditions<sup>18-21</sup>. These results suggest that SKPs have many characteristics that would be beneficial in nervous system regeneration<sup>22,23</sup>. In the previous study, however,



**Fig. 6.** A. Immunocytochemical studies for various neuronal and angiogenic marker proteins after the *in vitro* chemical neural induction of hSMSCs (scale bar=100 μm). Most marker proteins were highly visible 6 hrs after neural induction. Nestin was expressed in the 0 hr specimen (before nerve induction), which is similar to the result of FACS analysis. NGF and VEGF were highly visible with their receptors (p75NGFR, trkA, VEGFR1, and VEGFR2) during neuronal differentiation. B. Immunocytochemical intensities for specific proteins. The expression of most proteins, except NeuN, peaked 6 hrs after neural induction, and then decreased over time (24 hrs and 48 hrs after induction). Data represent the mean±SE of four independent experiments. A star (\*) indicates a significant difference from the control ( $P<0.05$ ). (S-100: S-100 protein, NF: neurofilament, MBP: myelin basic protein, NeuN: neural-specific nuclear protein, NGF: nerve growth factor, p75NGFR: p75 nerve growth factor receptor, trkA: tyrosine kinase receptor A, VEGF: vascular endothelial cell growth factor, VEGFR: vascular endothelial cell growth factor receptor)

Jun-Ho Byun et al: Isolation of human mesenchymal stem cells from the skin and their neurogenic differentiation *in vitro*. J Korean Assoc Oral Maxillofac Surg 2012



we isolated and cultured a different type of porcine skin-derived stem cells under culture condition different from that of SKPs as well as different serum-containing and adherent cell culturing method and observed the distinct MSC characteristics in these porcine skin-derived cells<sup>15</sup>. Similarly, in this study, human skin-derived cells were isolated from the submandibular skin segments, and multipotent MSC characteristics were detected, i.e., expression of transcription factors (Oct 4 and Nanog), detection of MSC markers (CD29, CD44, CD90, CD105, and vimentin), and potential of *in vitro* osteogenic, adipogenic, and chondrogenic differentiations. Interestingly, unlike the porcine SMSCs of a previous study<sup>15</sup>, we could not observe Sox 2 expression in the hSMSCs of this study. This is consistent with the result of other research, i.e., high visibility of Oct 4 and Nanog but negative expression of Sox 2 in human BMSCs<sup>24</sup>. In addition, they observed strongly expressed nestin, a neural precursor maker, in all kinds of MSCs originating in the bone marrow, dermis, and adipose tissues, similar to this study. Actually, this neural precursor marker was usually seen in SKPs, regarded as neural crest-originated stem cell in skin<sup>19</sup>. Taken together, the isolated and cultivated human skin-derived cells in this study are considered multipotent, hSMSCs demonstrating the characteristics of a neural precursor.

The *in vitro* neural differentiation methods of BMSCs have been investigated by several researchers. First, various molecules involved in neural development, such as growth factors, neurotrophins, cytokines, and retinoic acid, were used<sup>25,26</sup>. Second, neural differentiation was obtained by increasing the intracellular cyclic adenosine monophosphate<sup>27</sup>. Third, specific chemical compounds such as  $\beta$ -mercaptoethanol (BME), DMSO, and BHA were used in the serum-free culturing medium for neural induction<sup>16,17</sup>. Most recently, the neural differentiation of BMSCs was also observed when cultured in inflammatory astrocyte-containing medium<sup>28</sup>. In these various *in vitro* neural induction protocols, chemical protocols have been widely used and studied because it was a simple approach that rapidly yielded neuron-like cell differentiation from BMSCs<sup>29-31</sup>. Morphological changes into nerve-like cells and expression of neuron-specific makers have been observed within a few hours of chemical neural induction. In addition, more than 70% of differentiated neuron-like cells from BMSCs were observed after chemical neural induction, which was extremely higher concentration compared with the other induction method<sup>32,33</sup>.

In recent studies, however, the use of chemical compounds in serum-free conditions for the neural induction of BMSCs

produces toxic and stressful culture conditions. Therefore, these conditions contributed to the rapid disruption of the actin cytoskeleton and shape change to a form resembling nerve cells<sup>34,35</sup>. These nerve-like cells were not truly differentiated nerve cells, and they did not exhibit electrical conduction. Moreover, the chemical induction led to cell apoptosis and death<sup>33,36</sup>. In this study, we observed rapid neuron-like morphological changes of hSMSCs after chemical induction. After 6 hrs of induction, the highest neuron-like cells morphology and peak expression of neural and angiogenic-related proteins were detected. With the passage of induction time, however, the morphology of neural induced cells deteriorated, and the cell number and expression of nerve-related proteins decreased. Interestingly, when the induction medium was changed to DMEM with FBS, the neural differentiated cells returned to their original shape, and the cell number also increased. These results indicate that the chemically induced neuron-like cells of this study were not regarded as true nerve cells. When stem cells differentiate into target cells, it is difficult for them to return to stem cells especially in a short time period<sup>37</sup>. Therefore, the chemical induction protocol wherein the induction media contained BME, DMSO, and BHA was not a suitable choice for the neural differentiation method of hSMSCs, even though neuron-like morphology and various nerve-related proteins were observed shortly after induction.

In previous reports, SKPs, considered one of the neural crest originated-stem cells, were differentiated into nerve cells and maintained by supplementation with various growth factors and neurotrophins such as B27, NGF, brain-derived neurotrophic factor (BDNF), neuro-3 (NT-3), and FBS<sup>8,18,19</sup>. This is similar to previously reported neural differentiation methods of BMSCs involving various growth factors and neurotrophins<sup>25</sup>. Although further studies are needed to determine the most optimal method for *in vitro* neural differentiation from MSCs, the use of these growth factors and neurotrophins can be considered a substitute method for the *in vitro* neural induction of SMSCs.

Several researchers have demonstrated the enhanced nerve regeneration after the *in vivo* transplantation of differentiated Schwann cells or neurons from BMSCs<sup>4,5,38</sup>. Nonetheless, undifferentiated MSCs are well-known to have potential for differentiation into target cells after *in vivo* transplantation. Moreover, undifferentiated BMSCs expressed various neurotrophic factors after *in vivo* transplantation into the nerve defect site<sup>4</sup>. Since previous studies have not suggested an obvious *in vitro* neural differentiation method

from hSMSCs, undifferentiated autologous SMSCs could be suggested as reasonable transplant material for the regeneration of nerve defects.

## V. Conclusion

Isolated hSMSCs from adult skin showed MSC characteristics possessing multipotency and neural precursor marker, suggesting that human skin may be used as available adult stem cell source. Nonetheless, whether the *in vitro* neurogenic differentiated cells from hSMSCs under the chemical neural induction protocol produce actual nerve cells was doubtful, even though neuron-like cell morphology and various nerve-related proteins were detected in these chemically induced cells. Therefore, we suggest that the undifferentiated autologous hSMSCs - instead of chemically induced neuron-like cells - be considered for use as transplant material for *in vivo* nerve regeneration in the future.

## References

- Mosahebi A, Woodward B, Wiberg M, Martin R, Terenghi G. Retroviral labeling of Schwann cells: in vitro characterization and in vivo transplantation to improve peripheral nerve regeneration. *Glia* 2001;34:8-17.
- Murakami T, Fujimoto Y, Yasunaga Y, Ishida O, Tanaka N, Ikuta Y, et al. Transplanted neuronal progenitor cells in a peripheral nerve gap promote nerve repair. *Brain Res* 2003;974:17-24.
- Heine W, Conant K, Griffin JW, Höke A. Transplanted neural stem cells promote axonal regeneration through chronically denervated peripheral nerves. *Exp Neurol* 2004;189:231-40.
- Tohill M, Mantovani C, Wiberg M, Terenghi G. Rat bone marrow mesenchymal stem cells express glial markers and stimulate nerve regeneration. *Neurosci Lett* 2004;362:200-3.
- Dezawa M. Systematic neuronal and muscle induction systems in bone marrow stromal cells: the potential for tissue reconstruction in neurodegenerative and muscle degenerative diseases. *Med Mol Morphol* 2008;41:14-9.
- Cuevas P, Carceller F, Garcia-gómez I, Yan M, Dujovny M. Bone marrow stromal cell implantation for peripheral nerve repair. *Neurol Res* 2004;26:230-2.
- Hou SY, Zhang HY, Quan DP, Liu XL, Zhu JK. Tissue-engineered peripheral nerve grafting by differentiated bone marrow stromal cells. *Neuroscience* 2006;140:101-10.
- Toma JG, Akhavan M, Fernandes KJ, Barnabé-heider F, Sadikot A, Kaplan DR, et al. Isolation of multipotent adult stem cells from the dermis of mammalian skin. *Nat Cell Biol* 2001;3:778-84.
- Toma JG, McKenzie IA, Bagli D, Miller FD. Isolation and characterization of multipotent skin-derived precursors from human skin. *Stem Cells* 2005;23:727-37.
- Shi C, Zhu Y, Su Y, Cheng T. Stem cells and their applications in skin-cell therapy. *Trends Biotechnol* 2006;24:48-52.
- Dyce PW, Zhu H, Craig J, Li J. Stem cells with multilineage potential derived from porcine skin. *Biochem Biophys Res Commun* 2004;316:651-8.
- Blanpain C, Fuchs E. Epidermal stem cells of the skin. *Annu Rev Cell Dev Biol* 2006;22:339-73.
- Ohyama M, Terunuma A, Tock CL, Radonovich MF, Pise-masison CA, Hopping SB, et al. Characterization and isolation of stem cell-enriched human hair follicle bulge cells. *J Clin Invest* 2006;116:249-60.
- Riekstina U, Muceniece R, Cakstina I, Muiznieks I, Ancans J. Characterization of human skin-derived mesenchymal stem cell proliferation rate in different growth conditions. *Cytotechnology* 2008;58:153-62.
- Kang EJ, Byun JH, Choi YJ, Maeng GH, Lee SL, Kang DH, et al. In vitro and in vivo osteogenesis of porcine skin-derived mesenchymal stem cell-like cells with a demineralized bone and fibrin glue scaffold. *Tissue Eng Part A* 2010;16:815-27.
- Woodbury D, Schwarz EJ, Prockop DJ, Black IB. Adult rat and human bone marrow stromal cells differentiate into neurons. *J Neurosci Res* 2000;61:364-70.
- Woodbury D, Reynolds K, Black IB. Adult bone marrow stromal stem cells express germline, ectodermal, endodermal, and mesodermal genes prior to neurogenesis. *J Neurosci Res* 2002;69:908-17.
- Fernandes KJ, McKenzie IA, Mill P, Smith KM, Akhavan M, Barnabé-heider F, et al. A dermal niche for multipotent adult skin-derived precursor cells. *Nat Cell Biol* 2004;6:1082-93.
- Fernandes KJ, Kobayashi NR, Gallagher CJ, Barnabé-heider F, Aumont A, Kaplan DR, et al. Analysis of the neurogenic potential of multipotent skin-derived precursors. *Exp Neurol* 2006;201:32-48.
- Hunt DP, Morris PN, Sterling J, Anderson JA, Joannides A, Jahoda C, et al. A highly enriched niche of precursor cells with neuronal and glial potential within the hair follicle dermal papilla of adult skin. *Stem Cells* 2008;26:163-72.
- Zhao M, Isom SC, Lin H, Hao Y, Zhang Y, Zhao J, et al. Tracing the stemness of porcine skin-derived progenitors (pSKP) back to specific marker gene expression. *Cloning Stem Cells* 2009;11:111-22.
- Shih DT, Lee DC, Chen SC, Tsai RY, Huang CT, Tsai CC, et al. Isolation and characterization of neurogenic mesenchymal stem cells in human scalp tissue. *Stem Cells* 2005;23:1012-20.
- Marchesi C, Pluderi M, Colleoni F, Belicchi M, Merigalli M, Farini A, et al. Skin-derived stem cells transplanted into resorbable guides provide functional nerve regeneration after sciatic nerve resection. *Glia* 2007;55:425-38.
- Riekstina U, Cakstina I, Parfejevs V, Hoogduijn M, Jankovskis G, Muiznieks I, et al. Embryonic stem cell marker expression pattern in human mesenchymal stem cells derived from bone marrow, adipose tissue, heart and dermis. *Stem Cell Rev* 2009;5:378-86.
- Sanchez-ramos J, Song S, Cardozo-pelaez F, Hazzi C, Stedeford T, Willing A, et al. Adult bone marrow stromal cells differentiate into neural cells in vitro. *Exp Neurol* 2000;164:247-56.
- Kim BJ, Seo JH, Buben JK, Oh YS. Differentiation of adult bone marrow stem cells into neuroprogenitor cells in vitro. *Neuroreport* 2002;13:1185-8.
- Deng W, Obrocka M, Fischer I, Prockop DJ. In vitro differentiation of human marrow stromal cells into early progenitors of neural cells by conditions that increase intracellular cyclic AMP. *Biochem Biophys Res Commun* 2001;282:148-52.
- Wang FW, Jia DY, Du ZH, Fu J, Zhao SD, Liu SM, et al. Roles of activated astrocytes in bone marrow stromal cell proliferation and differentiation. *Neuroscience* 2009;160:319-29.
- Bossolasco P, Cova L, Calzarossa C, Rimoldi SG, Borsotti C, Deliliers GL, et al. Neuro-glial differentiation of human bone marrow stem cells in vitro. *Exp Neurol* 2005;193:312-25.
- Lei Z, Yongda L, Jun M, Yingyu S, Shaoju Z, Xinwen Z, et al. Culture and neural differentiation of rat bone marrow mesenchymal stem cells in vitro. *Cell Biol Int* 2007;31:916-23.
- Yamaguchi S, Kuroda S, Kobayashi H, Shichinohe H, Yano S, Hida K, et al. The effects of neuronal induction on gene expression profile in bone marrow stromal cells (BMSC)--a preliminary study using microarray analysis. *Brain Res* 2006;1087:15-27.

32. Muñoz-elías G, Woodbury D, Black IB. Marrow stromal cells, mitosis, and neuronal differentiation: stem cell and precursor functions. *Stem Cells* 2003;21:437-48.
33. Barnabé GF, Schwindt TT, Calcagnotto ME, Motta FL, Martinez GJ, De OA, et al. Chemically-induced RAT mesenchymal stem cells adopt molecular properties of neuronal-like cells but do not have basic neuronal functional properties. *PLoS One* 2009;4:e5222.
34. Lu P, Blesch A, Tuszynski MH. Induction of bone marrow stromal cells to neurons: differentiation, transdifferentiation, or artifact? *J Neurosci Res* 2004;77:174-91.
35. Neuhuber B, Gallo G, Howard L, Kostura L, Mackay A, Fischer I. Reevaluation of in vitro differentiation protocols for bone marrow stromal cells: disruption of actin cytoskeleton induces rapid morphological changes and mimics neuronal phenotype. *J Neurosci Res* 2004;77:192-204.
36. Rismanchi N, Floyd CL, Berman RF, Lyeth BG. Cell death and long-term maintenance of neuron-like state after differentiation of rat bone marrow stromal cells: a comparison of protocols. *Brain Res* 2003;991:46-55.
37. Rho GJ, Kumar BM, Balasubramanian SS. Porcine mesenchymal stem cells--current technological status and future perspective. *Front Biosci* 2009;14:3942-61.
38. Shimizu S, Kitada M, Ishikawa H, Itokazu Y, Wakao S, Dezawa M. Peripheral nerve regeneration by the in vitro differentiated-human bone marrow stromal cells with Schwann cell property. *Biochem Biophys Res Commun* 2007;359:915-20.