

Cryopreservation of dental tissue and subsequent isolation of mesenchymal stem cells

Bong-Wook Park, D.D.S., Ph.D.

Section Editor of JKAOMS

*Department of Oral and Maxillofacial Surgery, Institute of Health Sciences,
Gyeongsang National University School of Medicine, Jinju, Korea*

Mesenchymal stem cells (MSCs) are characterized as plate-adherent cells that demonstrate a fibroblast-like growth pattern, express specific cell surface markers (CD13, CD29, CD44, CD73, CD90, CD105, and CD166), and differentiate into mesenchymal lineage cells *in vitro* (osteocytes, chondrocytes, and adipocytes)¹. They were first isolated from bone marrow, but have since been isolated from various tissues, including skin, fat, cartilage, blood, umbilical cord, and dental tissue¹⁻³. Recently, dental tissue has emerged as a potential source of autologous MSCs that can be used for tissue regeneration via tissue engineering. Precursor cells in dental tissue were first isolated and characterized from dental pulp, after which cells showing the characteristics of MSCs were isolated from immature wisdom teeth, periodontal ligament, and exfoliated deciduous teeth³⁻⁶. In particular, the dental pulp, follicle, and root apical papilla of immature wisdom teeth have been reported to possess abundant undifferentiated primitive cells^{2,3,6}. Interestingly, stem cells from different dental tissues show similar MSC characteristics, but varying *in vitro* and *in vivo* differentiation properties^{2,3,6}. Stem cells from the dental follicle of extracted wisdom teeth exhibit superior osteogenic differentiation potential compared to those from the dental pulp or root apical papilla^{3,6}. In addition, one of the advantages of using dental stem cells from extracted wisdom teeth is that they can be obtained from patients at a relatively late age—from the late teens to mid-twenties. Moreover, recently, a new method for long-term cryopreservation of dental tissue from extracted wisdom teeth has been developed, and its efficacy was confirmed by the >70% cell survival rate of long-term preserved dental tissue⁷.

MSCs have potential clinical applications in the treatment of autoimmune diseases or reduction of immune responses after allotransplantation, as well as in tissue regeneration⁸. Many researchers have shown that MSCs exert immune suppression and anti-inflammatory activity by dominantly

inhibiting T-helper cells while activating regulatory T-cells and suppressing B-lymphocytes^{8,9}. This immunomodulatory effect of MSCs makes them suitable for treatment of severe autoimmune diseases and graft-versus-host disease⁸. However, most trials for the clinical application of MSCs have used bone marrow-derived MSCs (BMSCs). As mentioned previously, MSCs from dental tissue (DMSCs) exhibit similar characteristics to BMSCs, but have superior osteogenic differentiation potential^{3,6}. Therefore, DMSCs could possibly replace BMSCs in cell therapy and tissue regeneration applications. The new cryopreservation method for dental tissue enables the use of autologous MSCs from preserved dental tissue of extracted wisdom teeth⁷. These autologous MSCs could reduce unexpected side effects during clinical use, while maintaining similar immunomodulatory efficacy.

In conclusion, dental tissue from immature wisdom teeth, including that from the dental follicle, pulp, and root apical papilla, possesses abundant multipotent stem cells that could be useful in both cell therapy and tissue regeneration. Traditionally, these dental tissues have been discarded after extraction. In the future, dental tissue obtained via extraction of impacted teeth should be cryopreserved for use as an autologous stem cell source. The long-term preservation of dental tissue after tooth extraction could thus create a new business model in dentistry, as well as provide greater access to autologous MSCs for patients in need.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

References

1. Kang EJ, Byun JH, Choi YJ, Maeng GH, Lee SL, Kang DH, et al.

- In vitro and in vivo osteogenesis of porcine skin-derived mesenchymal stem cell-like cells with a demineralized bone and fibrin glue scaffold. *Tissue Eng Part A* 2010;16:815-27.
2. Song JH, Park BW, Byun JH, Kang EJ, Rho GJ, Shin SH, et al. Isolation and characterization of human dental tissue-derived stem cells in the impacted wisdom teeth: comparison of dental follicle, dental pulp, and root apical papilla-derived cells. *J Korean Assoc Oral Maxillofac Surg* 2010;36:186-96.
 3. Park BW, Kang EJ, Byun JH, Son MG, Kim HJ, Hah YS, et al. In vitro and in vivo osteogenesis of human mesenchymal stem cells derived from skin, bone marrow and dental follicle tissues. *Differentiation* 2012;83:249-59.
 4. Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci U S A* 2000;97:13625-30.
 5. Seo BM, Miura M, Gronthos S, Bartold PM, Batouli S, Brahimi J, et al. Investigation of multipotent postnatal stem cells from human periodontal ligament. *Lancet* 2004;364:149-55.
 6. Jeon BG, Kang EJ, Kumar BM, Maeng GH, Ock SA, Kwack DO, et al. Comparative analysis of telomere length, telomerase and reverse transcriptase activity in human dental stem cells. *Cell Transplant* 2011;20:1693-705.
 7. Park BW, Jang SJ, Byun JH, Kang YH, Choi MJ, Park WU, et al. Cryopreservation of human dental follicle tissue for use as a resource of autologous mesenchymal stem cells. *J Tissue Eng Regen Med* 2014. doi: 10.1002/term.1945. [Epub ahead of print]
 8. Zhao K, Lou R, Huang F, Peng Y, Jiang Z, Huang K, et al. Immunomodulation effects of mesenchymal stromal cells on acute graft-versus-host disease after hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant* 2015;21:97-104.
 9. Duffy MM, Ritter T, Ceredig R, Griffin MD. Mesenchymal stem cell effects on T-cell effector pathways. *Stem Cell Res Ther* 2011;2:34.